



Nuclear Structure for $0\nu\beta\beta$

Low Energy Community Meeting - 10 August 2020
FRIB-TA Annual Meeting

Saori Pastore
Washington University in St Louis

<https://physics.wustl.edu/quantum-monte-carlo-group>

Quantum Monte Carlo Group @ WashU

Lorenzo Andreoli (PD) Jason Bub (GS) Garrett King (GS) Maria Piarulli and Saori Pastore

Computational Resources awarded by the DOE ALCC and INCITE programs

Nuclei for Fundamental Symmetries & Neutrinos


Nuclei are used for precision tests of the standard model and in searches for physics beyond the standard model.

An accurate understanding of nuclear structure and dynamics in a wide range of energy and momentum transferred is required in order to disentangle new physics from nuclear effects.


In this talk, I will focus on the **role of two- and three-body correlations and currents in selected nuclear electroweak observables** at different kinematics.

The emphasis will be on light nuclei ($A \leq 12$). For these systems Variational and Green's Function Monte Carlo methods allow to retain many-body effects and provide results with a computational accuracy (in most cases) of the order of few percents.

Ground States'
Electroweak Moments,
Form Factors, Radii



Neutrinoless Double
Beta Decay,
Muon-Capture



Accelerator Neutrino
Experiments,
Lepton-Nucleus XSecs

$(\omega, q) \sim 0$ MeV

$\omega \sim \text{few MeVs}$
 $q \sim 0$ MeV

$\omega \sim \text{few MeVs}$
 $q \sim 10^2$ MeV

$\omega \sim \text{tens of MeVs}$

$\omega \sim 10^2$ MeV



Electromagnetic
Decay, Beta Decay,
Double Beta Decay &
inverse processes



Nuclear Rates for
Astrophysics



Strategy

Validate the Nuclear Model against available data for strong and electroweak observables

- Energy Spectra, Electromagnetic Form Factors, Electromagnetic Moments, ...
- Electromagnetic and **Beta decay rates**, ...
- Muon Capture Rates, ...
- **Electron-Nucleus Scattering Cross Sections**, ...

Use attained information to make (accurate) predictions for BSM searches and precision tests


- EDMs, Anapole Moments, Hadronic PV, ...
- BSM searches with beta decay, ...
- **Neutrinoless double beta decay**, ...
- Neutrino-Nucleus Scattering Cross Sections, ...
- ...

Many-body Nuclear Interactions

Many-body Nuclear Hamiltonian

$$H = T + V = \sum_{i=1}^A t_i + \sum_{i<j} v_{ij} + \sum_{i<j<k} V_{ijk} + \dots$$


v_{ij} and V_{ijk} are two- and three-nucleon operators based on experimental data fitting; fitted parameters subsume underlying QCD dynamics



Contact term: short-range

Two-pion range: intermediate-range $r \propto (2m_\pi)^{-1}$

One-pion range: long-range $r \propto m_\pi^{-1}$



SP et al. PRC80(2009)034004


In Quantum Monte Carlo methods we use:

AV18+UIX; **AV18+IL7** Wiringa, Schiavilla, Pieper *et al.*


chiral π N **N2LO+N2LO** Gerzelis, Tews, Lynn *et al.*

chiral π N Δ **N3LO+N2LO** Piarulli *et al.* **Norfolk Models**

Many-body Nuclear Electroweak Currents



one-body



two-body

- One-body currents: non-relativistic reduction of covariant nucleons' currents
- Two-body currents are a manifestation of two-nucleon correlations
- Electromagnetic two-body currents are required to satisfy current conservation


$$\mathbf{q} \cdot \mathbf{j} = [H, \rho] = [t_i + v_{ij} + V_{ijk}, \rho]$$

Nuclear Charge Operator

$$\rho = \sum_{i=1}^A \rho_i + \sum_{i<j} \rho_{ij} + \dots$$

Nuclear (Vector) Current Operator


$$\mathbf{j} = \sum_{i=1}^A \mathbf{j}_i + \sum_{i<j} \mathbf{j}_{ij} + \dots$$



Magnetic Moment: Single Particle Picture


Neutrino-Nucleus Interactions

Low energy and momentum:
Beta Decay Matrix Elements




Gysbers *et al.* NaturePhys15(2019)

High Energy (on Nuclear Physics Scale):
Neutrino-Nucleus Cross Section



Alvarez-Ruso arXiv:1012.3871

Beta Decay and Electron Capture in Light Nuclei




Calculations based on



- chiral interactions and currents
NV2+3-Ia Norfolk unstarred
NV2+3-Ia* Norfolk* starred
 Piarulli *et al.* PRL120(2018)052503
 Baroni *et al.* PRC98(2018)044003
- phenomenological **AV18+IL7**
 potential and chiral axial currents
 (hybrid calculation)

Two-body currents are small/negligible;
 Results for $A=6-7$ are within 2% of data;
 Results for $A=8$ are off by a 30-40%;
 Results for $A=10$ are affected by the
 second $J^\pi=(1^+)$ state in ${}^{10}\text{B}$

Three-body Force and the Axial Contact Current



Three-body force




Axial two-body contact current

LECs c_D and c_E are fitted to:

- trinucleon B.E. and nd doublet scattering length in **NV2+3-Ia**
- trinucleon B.E. and Gamow-Teller matrix element of tritium **NV2+3-Ia***

Baroni *et al.* PRC98(2018)044003

Axial Two-body Transition Density



NV2+3-la ; NV2+3-la*

enhanced contribution from contact current in the starred model gives rise to nodes in the two-body transition density

Two-body axial currents




long-range at N2LO and N3LO



contact current at N3LO


Scaling and Universality of Short-Range Dynamics



G. King *et al.* to appear in PRC(2020) arXiv:2004.05263

NV2+3-Ia empty circles; NV2+3-Ia* stars
Different colors refer to different transitions

Neutrinoless Double Beta Decay



Nuclear matrix elements for neutrinoless double beta decay are required to extract the neutrino parameters from the decay rate (if the process is observed)

Matrix elements for nuclei of experimental interest are currently affected by large uncertainties due to truncation in the model space and partial (or missing) inclusion of many-body effects


We study neutrinoless double beta decay in **light nuclei** that have been successfully described by *ab initio* models where correlations and currents can be fully accounted for

These studies serve as benchmark and to establish the relevance of the various two-body (or more) dynamics inducing the decay

Engel & Menéndez

Rep.Progr.Phys80(2017)046301

Neutrinoless Double Beta Decay Matrix Elements




SP *et al.* PRC97(2018)014606



Cirigliano Dekens DeVries Graesser Mereghetti *et al.*
 PLB769(2017)460, JHEP12(2017)082, PRC97(2018)065501

- Leading operators in neutrinoless double beta decay are two-body operators
- These observables are particularly sensitive to short-range and two-body physics
- Transition densities calculated in momentum space indicate that the momentum transfer in this process is of the order of ~ 200 MeV

Comparison with Shell-Model Calculations



Closer agreement between Shell-Model calculations with Variational Monte Carlo results is reached by

- Increasing the size of the model space
- Wood-Saxon single particle wave functions are superior in describing the tails of the densities wrt harmonic oscillator wave functions
- Phenomenological Short-Range-Correlations functions further improve the agreement

Lepton-Nucleus scattering: Inclusive Processes

Electromagnetic Nuclear Response Functions

$$R_{\alpha}(q, \omega) = \sum_f \delta(\omega + E_0 - E_f) |\langle f | O_{\alpha}(\mathbf{q}) | 0 \rangle|^2$$

Longitudinal response induced by the charge operator $O_L = \rho$

Transverse response induced by the current operator $O_T = \mathbf{j}$

5 Responses in neutrino-nucleus scattering

The Quantum Monte Carlo community at LANL, ANL, JLAB delivered calculations of inclusive responses of both electron and neutrino scattering from nuclei with mass number $A \leq 12$


For a recent review see Rocco *Front.inPhys.*8 (2020)116

Lepton-Nucleus scattering: Data



Transverse Sum Rule

$$S_T(q) \propto \langle 0 | \mathbf{j}^\dagger \mathbf{j} | 0 \rangle \propto \langle 0 | \mathbf{j}_{1b}^\dagger \mathbf{j}_{1b} | 0 \rangle + \langle 0 | \mathbf{j}_{1b}^\dagger \mathbf{j}_{2b} | 0 \rangle + \dots$$

Observed transverse enhancement explained by the combined effect of two-body correlations and currents in the interference term



^{12}C Electromagnetic Data
Benhar *et al.* RMP80(2008)198





Transverse/Longitudinal Sum Rule
Carlson *et al.* PRC65(2002)024002

Beyond Inclusive: Short-Time-Approximation


Short-Time-Approximation Goals:

- Describe electroweak scattering from $A > 12$ without losing two-body physics
- Account for exclusive processes
- Incorporate relativistic effects



[Stanford Lab article](#)

Short-Time-Approximation



Short-Time-Approximation:

- Based on Factorization
- Allows to retain both two-body correlations and currents at the vertex
- Provides “more” exclusive information in terms of nucleon-pair kinematics via the Response Densities
- Response functions are given by the scattering from pairs of fully interacting nucleons that propagate into a correlated pair of nucleons
- Correctly accounts for interference

Response Functions

$$R_{\alpha}(q, \omega) = \sum_f \delta(\omega + E_0 - E_f) |\langle f | O_{\alpha}(\mathbf{q}) | 0 \rangle|^2$$


Response Densities

$$R(q, \omega) \sim \int \delta(\omega + E_0 - E_f) dP' dp' \mathcal{D}(p', P'; q)$$


P' and p' are the CM and relative momenta of the struck nucleon pair

Transverse Response Density: e - ${}^4\text{He}$ scattering

Transverse Density $q = 500 \text{ MeV}/c$




$e^{-4}\text{He}$ scattering in the back-to-back kinematic



SP *et al.* PRC101(2020)044612

- pp pairs
- nn pairs
- all pairs 1body
- all pairs tot



Subedi *et al.* Science320(2008)1475

Summary and Outlook

Ab initio calculations of light nuclei yield a picture of nuclear structure and dynamics where many-body effects play an essential role to explain available data.

This is found in a wide range of energy and momentum transferred and in both static and dynamical observables.

Light to low and medium mass nuclei are promising candidates for precision tests and studies of fundamental symmetries and neutrino physics.

In these systems, the existing computational methods allow to retain the complexity of many-body correlations and currents.

The QMC community is addressing larger nuclear systems, developing new algorithms that allow to retain two-body physics (correlations and currents) in lepton-nucleus scattering, and studying observables in kinematic regions of interest to experimental programs in BSM and precision physics.

Strategy

Validate the Nuclear Model against available data for strong and electroweak observables

- Energy Spectra, Electromagnetic Form Factors, Electromagnetic Moments, ...
- Electromagnetic and Beta decay rates, ...
- Muon Capture Rates, ...
- Electron-Nucleus Scattering Cross Sections, ...

Use attained information to make (accurate) predictions for BSM searches and precision tests

- EDMs, Anapole Moments, Hadronic PV, ...
- BSM searches with beta decay, ...
- Neutrinoless double beta decay, ...
- Neutrino-Nucleus Scattering Cross Sections, ...
- ...

Collaborators

WashU: **Andreoli Bub King** Piarulli

LANL: **Baroni** Carlson Cirigliano Gandolfi Hayes Mereghetti

JLab+ODU: Schiavilla

ANL: Lovato Rocco Wiringa

UCSD/UW: Dekens

Pisa U/INFN: Kievsky Marcucci Viviani

Salento U: Girlanda


Huzhou U: Dong Wang




Theory Alliance
FACILITY FOR RARE ISOTOPE BEAMS



Ground States'
Electroweak Moments,
Form Factors, Radii



Neutrinoless Double
Beta Decay,
Muon-Capture



Accelerator Neutrino
Experiments,
Lepton-Nucleus XSecs

$(\omega, q) \sim 0$ MeV

$\omega \sim \text{few MeVs}$
 $q \sim 0$ MeV

$\omega \sim \text{few MeVs}$
 $q \sim 10^2$ MeV

$\omega \sim \text{tens of MeVs}$

$\omega \sim 10^2$ MeV



Electromagnetic
Decay, Beta Decay,
Double Beta Decay &
inverse processes



Nuclear Rates for
Astrophysics

